

# LOGARITHMIC CONVEXITY OF GINI MEANS

FENG QI AND BAI-NI GUO

ABSTRACT. In the paper, the monotonicity and logarithmic convexity of Gini means and related functions are investigated.

## 1. INTRODUCTION

Recall [7] that Gini means were defined as

$$G(r, s; x, y) = \begin{cases} \left( \frac{x^s + y^s}{x^r + y^r} \right)^{1/(s-r)}, & r \neq s; \\ \exp\left( \frac{x^r \ln x + y^r \ln y}{x^r + y^r} \right), & r = s; \end{cases} \quad (1)$$

where  $x$  and  $y$  are positive variables and  $r$  and  $s$  are real variables. They are also called sum mean values.

There has been a lot of literature such as [3, 4, 5, 6, 9, 10, 11, 12, 13, 19, 20, 21] and the related references therein about inequalities and properties of Gini means.

The aim of this paper is to prove the monotonicity and logarithmic convexity of Gini means  $G(r, s; x, y)$  and related functions.

**Theorem 1.** *Gini means  $G(r, s; x, y)$  are*

- (1) *increasing with respect to both  $r \in (-\infty, \infty)$  and  $s \in (-\infty, \infty)$ ;*
- (2) *logarithmically convex with respect to both  $r$  and  $s$  if  $(r, s) \in (-\infty, 0) \times (-\infty, 0)$ ;*
- (3) *logarithmically concave with respect to both  $r$  and  $s$  if  $(r, s) \in (-\infty, 0) \times (-\infty, 0)$ .*

**Theorem 2.** *Let*

$$H_{r,s;x,y}(t) = G(r+t, s+t; x, y) \quad (2)$$

*for  $t \in \mathbb{R}$ . Then Gini means  $H_{r,s;x,y}(t)$  are*

- (1) *increasing on  $(-\infty, \infty)$ ;*
- (2) *logarithmically convex on  $(-\infty, -\frac{r+s}{2})$ ;*
- (3) *logarithmically concave on  $(-\frac{r+s}{2}, \infty)$*

*and the function*

$$K_{r,s;x,y}(t) = H_{r,s;x,y}(t)H_{r,s;x,y}(-t) \quad (3)$$

*is*

- (1) *increasing on  $(-\infty, 0)$ ;*
- (2) *decreasing on  $(0, \infty)$ .*

**Theorem 3.** *The function  $t \mapsto t \ln H_{r,s;x,y}(t)$  is convex*

- (1) *on  $(-\frac{s+r}{2}, 0)$  if  $s+r > 0$ ;*
- (2) *on  $(0, -\frac{s+r}{2})$  if  $s+r < 0$ .*

2000 *Mathematics Subject Classification.* Primary 26A48, 26A51; Secondary 26B25, 26D07.

*Key words and phrases.* Gini means, logarithmic convexity, monotonicity.

The first author was partially supported by the China Scholarship Council.

This paper was typeset using  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$ .

*Remark 1.* The extended mean values  $E(r, s; x, y)$  have properties similar to those obtained in the above theorems, see [1, 2, 8, 16, 17]. Similar problems were also discussed in [14, 15].

*Remark 2.* For completeness, although the monotonicity of Gini means  $G(r, s; x, y)$  has been verified in [6, 7] and related references, we would also like to give it a proof in the next section.

## 2. PROOFS OF THEOREMS

*Proof of Theorem 1.* It is easy to see that

$$\ln G(r, s; x, y) = \begin{cases} \frac{1}{s-r} \int_r^s \frac{x^u \ln x + y^u \ln y}{x^u + y^u} du, & r \neq s, \\ \frac{x^r \ln x + y^r \ln y}{x^r + y^r}, & r = s. \end{cases} \quad (4)$$

Since

$$\frac{d}{du} \left[ \frac{x^u \ln x + y^u \ln y}{x^u + y^u} \right] = \frac{x^u y^u (\ln x - \ln y)^2}{(x^u + y^u)^2} > 0 \quad (5)$$

and

$$\frac{d^2}{du^2} \left[ \frac{x^u \ln x + y^u \ln y}{x^u + y^u} \right] = -\frac{x^u y^u (x^u - y^u) (\ln x - \ln y)^3}{(x^u + y^u)^3} = \begin{cases} \geq 0, & u \leq 0, \\ \leq 0, & u \geq 0, \end{cases} \quad (6)$$

then the integrand in (4) is increasing on  $u \in (-\infty, \infty)$ , logarithmically convex on  $u \in (-\infty, 0)$  and logarithmically concave on  $u \in (0, \infty)$ . It is known [18, Lemma 1] that if  $f(t)$  is differentiable and increasing on an interval  $I$ , then the integral arithmetic mean of  $f(t)$ ,

$$\phi(r, s) = \begin{cases} \frac{1}{s-r} \int_r^s f(t) dt, & r \neq s, \\ f(r), & r = s, \end{cases} \quad (7)$$

is also increasing with  $r$  and  $s$  on  $I$ ; If  $f(t)$  is twice differentiable and convex on  $I$ , then  $\phi(r, s)$  is also convex with  $r$  and  $s$  on  $I$ . Consequently, Gini means  $G(r, s; x, y)$  with respect to both  $r$  and  $s$  are increasing on  $(-\infty, \infty)$ , logarithmically convex if  $(r, s) \in (-\infty, 0) \times (-\infty, 0)$ , and logarithmically concave if  $(r, s) \in (0, \infty) \times (0, \infty)$ . The proof of Theorem 1 is complete.  $\square$

*Proof of Theorem 2.* Taking the logarithm of  $H_{r,s;x,y}(t)$  and differentiating consecutively yields

$$\begin{aligned} \ln H_{r,s;x,y}(t) &= \frac{1}{s-r} [\ln(x^{s+t} + y^{s+t}) - \ln(x^{r+t} + y^{r+t})], \\ [\ln H_{r,s;x,y}(t)]' &= \frac{1}{s-r} \left( \frac{x^{s+t} \ln x + y^{s+t} \ln y}{x^{s+t} + y^{s+t}} - \frac{x^{r+t} \ln x + y^{r+t} \ln y}{x^{r+t} + y^{r+t}} \right), \\ [\ln H_{r,s;x,y}(t)]'' &= \frac{1}{s-r} \left[ \frac{x^{s+t} y^{s+t} (\ln x - \ln y)^2}{(x^{s+t} + y^{s+t})^2} - \frac{x^{r+t} y^{r+t} (\ln x - \ln y)^2}{(x^{r+t} + y^{r+t})^2} \right]. \end{aligned}$$

By virtue of (5), it easily follows that  $[\ln H_{r,s;x,y}(t)]' \geq 0$ , which means that Gini means  $H_{r,s;x,y}(t)$  is increasing on  $(-\infty, \infty)$ .

With the aid of (6), it may be obtained easily that the function

$$f_{x,y}(u) = \frac{x^u y^u (\ln x - \ln y)^2}{(x^u + y^u)^2} \quad (8)$$

is increasing on  $u \in (-\infty, 0)$  and decreasing on  $(0, \infty)$ . Moreover, it is clear that  $f_{x,y}(u) = f_{x,y}(-u)$ , that is, the function  $f_{x,y}(u)$  is even on  $(-\infty, \infty)$ .

Let

$$F_{x,y}(t) = f_{x,y}(s+t) - f_{x,y}(r+t). \quad (9)$$

If  $s+t > r+t > 0$ , that is,  $t > -r > -s$ , since  $f_{x,y}(u)$  is decreasing on  $(0, \infty)$ , then  $F_{x,y}(t) \leq 0$ . Similarly, if  $r+t < s+t < 0$ , i.e.,  $t < -s < -r$ , then  $F_{x,y}(t) \geq 0$ . If  $r+t < 0 < s+t$  and  $0 < -(r+t) < s+t$ , equivalently,  $t > -\frac{r+s}{2}$ , since  $f_{x,y}(u)$  is even on  $(-\infty, \infty)$  and decreasing on  $(0, \infty)$ , then  $F_{x,y}(t) \leq 0$ ; Similarly, if  $t < -\frac{r+s}{2}$ , then  $F_{x,y}(t) \geq 0$ . This implies

$$[\ln H_{r,s;x,y}(t)]'' \begin{cases} \geq 0, & t < -\frac{r+s}{2} \\ \leq 0, & t > -\frac{r+s}{2} \end{cases}$$

for all  $r, s, x, y$  by a recourse to symmetric properties  $G(r, s; x, y) = G(s, r; x, y) = G(r, s; y, x)$ .

Taking the logarithm on both sides of (3) and differentiating gives

$$[\ln K_{r,s;x,y}(t)]' = \frac{H'_{r,s;x,y}(t)}{H_{r,s;x,y}(t)} - \frac{H'_{r,s;x,y}(-t)}{H_{r,s;x,y}(-t)},$$

where

$$H'_{r,s;x,y}(-t) = \left. \frac{dH_{r,s;x,y}(u)}{du} \right|_{u=-t}.$$

The logarithmic convexities of  $H_{r,s;x,y}(t)$  implies that the function  $\frac{H'_{r,s;x,y}(t)}{H_{r,s;x,y}(t)}$  is increasing on  $(-\infty, -\frac{r+s}{2})$  and decreasing on  $(-\frac{r+s}{2}, 0)$ . Careful computation can verify that

$$\frac{H'_{r,s;x,y}(t)}{H_{r,s;x,y}(t)} = \frac{H'_{r,s;x,y}(-t - (s+r))}{H_{r,s;x,y}(-t - (s+r))}$$

for  $t \in (-\infty, \infty)$ . Consequently, the function

$$Q(t) = \frac{H'_{r,s;x,y}(t - (s+r)/2)}{H_{r,s;x,y}(t - (s+r)/2)}$$

is increasing on  $(-\infty, 0)$  and decreasing on  $(0, \infty)$  and satisfies  $Q(t) = Q(-t)$  for  $t \in (-\infty, \infty)$ . Utilization of the approach applied to the function  $f_{x,y}$  above yields that  $Q(t + (s+r)) - Q(t)$  is positive on  $(-\infty, -\frac{s+r}{2})$  and negative on  $(-\frac{s+r}{2}, \infty)$ , which is equivalent to

$$Q\left(t + \frac{s+r}{2}\right) - Q\left(t - \frac{s+r}{2}\right) = \frac{H'_{r,s;x,y}(t)}{H_{r,s;x,y}(t)} - \frac{H'_{r,s;x,y}(t - (s+r))}{H_{r,s;x,y}(t - (s+r))} \quad (10)$$

being positive on  $(-\infty, 0)$  and negative on  $(0, \infty)$ . Since

$$K_{r,s;x,y}(t) = \frac{xyH_{r,s;x,y}(t)}{H_{r,s;x,y}(t - (s+r))}, \quad (11)$$

then the function in (10) equals  $[\ln K_{r,s;x,y}(t)]'$ , which implies that the function  $K_{r,s;x,y}(t)$  is increasing on  $(-\infty, 0)$  and decreasing on  $(0, \infty)$ . The proof of Theorem 2 is complete.  $\square$

*Proof of Theorem 3.* Direct calculation yields

$$[t \ln H_{r,s;x,y}(t)]'' = 2[\ln H_{r,s;x,y}(t)]' + t[\ln H_{r,s;x,y}(t)]''. \quad (12)$$

By Theorem 2, it follows that  $[\ln H_{r,s;x,y}(t)]' > 0$  on  $(-\infty, \infty)$ ,  $[\ln H_{r,s;x,y}(t)]'' > 0$  on  $(-\infty, -\frac{s+r}{2})$  and  $[\ln H_{r,s;x,y}(t)]'' < 0$  on  $(-\frac{s+r}{2}, \infty)$ . Therefore, if  $s+r < 0$  then  $[t \ln H_{r,s;x,y}(t)]'' > 0$  and  $t \ln H_{r,s;x,y}(t)$  is convex on  $(0, -\frac{s+r}{2})$ , if  $s+r > 0$

then  $[t \ln H_{r,s;x,y}(t)]'' > 0$  and  $t \ln H_{r,s;x,y}(t)$  is convex on  $(-\frac{s+r}{2}, 0)$ . The proof of Theorem 3 is complete.  $\square$

## REFERENCES

- [1] W.-S. Cheung and F. Qi, *Logarithmic convexity of the one-parameter mean values*, RGMIA Res. Rep. Coll. **7** (2004), no. 2, Art. 15, 331–342; Available online at <http://www.staff.vu.edu.au/rgmia/v7n2.asp>.
- [2] W.-S. Cheung and F. Qi, *Logarithmic convexity of the one-parameter mean values*, Taiwanese J. Math. **11** (2007), no. 1, 231–237.
- [3] P. Czinder and Zs. Páles, *A general Minkowski-type inequality for two variable Gini means*, Publ. Math. Debrecen **57** (2000), no. 1-2, 203–216.
- [4] P. Czinder and Zs. Páles, *Local monotonicity properties of two-variable Gini means and the comparison theorem revisited*, J. Math. Anal. Appl. **301** (2005), no. 2, 427–438.
- [5] Z. Daróczy and L. Losonczi, *Über den Vergleich von Mittelwerten*, Publ. Math. Debrecen **17** (1970), 289–297.
- [6] D. Farnsworth and R. Orr, *Gini means*, Amer. Math. Monthly **93** (1986), no. 8, 603–607.
- [7] C. Gini, *Di una formula compresive delle medie*, Metron **13** (1938), 3–22.
- [8] B.-N. Guo and F. Qi, *A simple proof of logarithmic convexity of extended mean values*, Numer. Algorithms (2009), in press; Available online at <http://dx.doi.org/10.1007/s11075-008-9259-7>.
- [9] Zh. Liu, *Minkowski's inequality for extended mean values*, Proceedings of the Second ISAAC Congress, Vol. 1 (Fukuoka, 1999), 585–592, Int. Soc. Anal. Appl. Comput., 7, Kluwer Acad. Publ., Dordrecht, 2000.
- [10] L. Losonczi, *Inequalities for integral mean values*, J. Math. Anal. Appl. **61** (1977), 586–606.
- [11] E. Neuman and Zs. Páles, *On comparison of Stolarsky and Gini means*, J. Math. Anal. Appl. **278** (2003), no. 2, 274–284.
- [12] E. Neuman and J. Sándor, *Inequalities involving Stolarsky and Gini means*, Math. Pannon. **14** (2003), no. 1, 29–44.
- [13] Zs. Páles, *Comparison of two variable homogeneous means*, General Inequalities 6. Proc. 6th Internat. Conf. Math. Res. Inst. Oberwolfach, Birkhäuser Verlag Basel, 1992, pp. 59–69.
- [14] F. Qi, *Complete monotonicity of logarithmic mean*, RGMIA Res. Rep. Coll. **10** (2007), no. 1, Art. 18; Available online at <http://www.staff.vu.edu.au/rgmia/v10n1.asp>.
- [15] F. Qi and Sh.-X. Chen, *Complete monotonicity of the logarithmic mean*, Math. Inequal. Appl. **10** (2007), no. 4, 799–804.
- [16] F. Qi, P. Cerone, S. S. Dragomir and H. M. Srivastava, *Alternative proofs for monotonic and logarithmically convex properties of one-parameter mean values*, Appl. Math. Comput. **208** (2009), no. 1, 129–133; Available online at <http://dx.doi.org/10.1016/j.amc.2008.11.023>.
- [17] F. Qi and B.-N. Guo, *The function  $(b^x - a^x)/x$ : Logarithmic convexity and applications to extended mean values*, Available online at <http://arxiv.org/abs/0903.1203>.
- [18] F. Qi, S.-L. Xu and L. Debnath, *A new proof of monotonicity for extended mean values*, Internat. J. Math. Math. Sci. **22** (1999), no. 2, 415–420.
- [19] J. Sándor, *A note on the Gini means*, Gen. Math. **12** (2004), no. 4, 17–21.
- [20] J. Sándor, *The Schur-convexity of Stolarsky and Gini means*, Banach J. Math. Anal. **1** (2007), no. 2, 212–215.
- [21] S. Toader and G. Toader, *Complementaries of Greek means with respect to Gini means*, Int. J. Appl. Math. Stat. **11** (2007), no. 7, 187–192.

(F. Qi) RESEARCH INSTITUTE OF MATHEMATICAL INEQUALITY THEORY, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

E-mail address: [qifeng618@gmail.com](mailto:qifeng618@gmail.com), [qifeng618@hotmail.com](mailto:qifeng618@hotmail.com), [qifeng618@qq.com](mailto:qifeng618@qq.com)

URL: <http://qifeng618.spaces.live.com>

(B.-N. Guo) SCHOOL OF MATHEMATICS AND INFORMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

E-mail address: [bai.ni.guo@gmail.com](mailto:bai.ni.guo@gmail.com), [bai.ni.guo@hotmail.com](mailto:bai.ni.guo@hotmail.com)

URL: <http://guobaini.spaces.live.com>